DOI: https://doi.org/10.67897/njma.2022.4y6be108

Nigerian Journal of Mathematics and Applications
Volume **32** (**II**), (2022), 151 – 168. *Printed by Unilorin press*
©*Nig. J. Math. Appl.* *http* : *//www.njmaman.com*

Special Issue in Honor of Prof. J. A. Gbadeyan's Retirement

Optimization of two-step method with Bhaskara points for solving non-linear dynamical problems

Umaru Mohammed[1*], Oyewole Abiodun Oyelami[2], Jamiu Garba[3], Abdullahi Aliyu Alhaji[4] and Mikhail Semenov[5]

Abstract

We derive a two-step method for solving non-linear dynamical problems employing Bhaskara points as off grid points. Deterministic in nature, non-linear dynamical systems display a periodic behavior that is highly dependent on the beginning circumstances, making long-term predictions impossible. Four Bhaskara points are generated as hybrid points to optimized the system. The method is use to solve non-linear dynamical systems to demonstrate the accuracy and efficient of the method. The method is zero stable and consistent. Some examples from literature were considered to demonstrate the efficiency and the accuracy of the method.

1. Introduction

Ordinary differential equations (ODEs) arise science and technology, whose solutions are not only important but also necessary and compulsory for the advancement of science and technology. Obtaining solutions to ODEs is as important as modelling that gives rise to them.

A linear ODE is one that can be split down into components, solved individually, and then recombined to produce the result. Nonlinear interactions take place whenever components of a system interact, compete, or cooperate. This kind of system is deterministic, but its long-term predictions are erroneous because of the periodic behavior it displays in response to the starting circumstances. However, most non-linear dynamical systems do not have analytical solutions, thereby relying on numerical methods. On of such methods

Received: 24/06/2022, Accepted: 28/07/2022, Revised: 02/08/2022. * Corresponding author.
2015 *Mathematics Subject Classification.* 65L07 & 35G99.
Keywords and phrases. collocation, interpolation, Bhaskara points, two-step method
[1&3]Department of Mathematics, Federal University of Technology, Minna, Nigeria
[2]Department of Mathematics, National Open University of Nigeria
[4]Department of Mechanical Engineering, Federal University of Technology, Minna, Nigeria
[5]Division for Experimental Physics, Tomsk Polytechnics University, Russia
E-mails of the corresponding author: umaru.mohd@futminna.edu.ng
ORCID of the corresponding author: 0000-0002-0777-6536

is the block method. Block methods were introduce by Milne [13]. The block methods proposed by Milne [13] has some draw backs, which led to the introduction of hybrid method. Hybrid methods has the advantage of easy change of step size and making use of data from off-set points that generally enhance the accuracy and effectiveness of the methods. The solution of non-linear dynamical problems is considered in this paper. Motsa [16] considered one-step solution of first order ODEs with equal space interval as hybrid point. Bothayna [10], considered single step method with equally three intra-step points for solving first order ODEs. Many researchers recently have applied different steps with different hybrid points for the development of numerical integrators for first order differential equation [3–7,14,20–23]. Recently, there have been discussions on how to generate intra-step points methodically, in order to yield optimal accuracy for block hybrid algorithms. By minimizing the local truncation errors (LTEs), Ramos [17] presented an optimized two-step block hybrid method for solving general first order IVPs. In [18], Ramos et al. expanded the concept of minimizing the LTE to derive optimized two-step block hybrid methods for the numerical solution of general second order IVPs. It is important to note that, in both [17] and [18], only two intra-step points were established. In light of the above we present a more direct method for the solution of first order non-linear ODEs, which will be applicable to special, stiff, non-linear and general forms of first order differential equations. The proposed method is a two-step hybrid method using four generated Bhaskara points as hybrid points to optimize the method. The proposed methods will be time efficient, have wider integration range and economically reliable. The aim of the study is to develop an optimized multi-derivative method with intra-step point for solving non-linear dynamical system and Partial Differential Equations.

We implement an implicit two-step method using Bhaskara points as hybrid points. The Gauss-Lobatto grid points are points generated from the algorithm $\{x\}_{i=0}^{M} = \frac{k}{2}\cos\frac{\pi i}{M} + \frac{k}{2}$, while Bhaskara points are points generated from approximating the cosine function $\{\cos\frac{\pi i}{M}\}_{i=0}^{M} \approx \frac{M^2-4i^2}{M^2+i^2}$ where $M = m+1$ is number of intra-steps and $M \in \mathbf{N} : m \geq 2$, and k is the step size.

We studied the order, zero stability, convergence and consistency of the method. Some numerical problem which are non-linear will be solved and compared to others in literature to show the efficiency and accuracy. The solution of first order initial value problem is of the form.

$$y' = f(x,y), \quad y(x_0) = y_0, \quad x_0 \in [a,b] \tag{1.1}$$

The function $f(x,y)$ satisfies the existence and uniqueness theorem and is continuous [8]. The hybrid block technique has recently been extended by a number of authors [3–7, 14, 20–23] to numerically solve first order differential equations with varied numbers of steps and hybrids.

2. Derivation of the Method

To derive the method, the power series of the form below is considered:

$$y(x) = \sum_{j=0}^{I+C_1+C_2} \alpha_j x^j \tag{2.1}$$

where I is the interpolation point, C_1 is the number of collocation points for the first derivative and C_2 is the number of collocation points for the second derivative as an approximate solution to the general first order problems of the form:

$$y' = f(x, y), \quad y(a) = \alpha. \tag{2.2}$$

The first and second derivatives of (2.1) are

$$y'(x) = \sum_{j=1}^{I+C_1+C_2} j\alpha_j x^{j-1} \tag{2.3}$$

$$y''(x) = \sum_{j=2}^{I+C_1+C_2} j(j-1)\alpha_j x^{j-2} \tag{2.4}$$

(2.1) is interpolated at $x = x_{n+1}$ while (2.3) and (2.4) are collocated at

$$x = x_{n+i}, \quad i = 0, 5/26, 20/29, 1, 38/29, 47/26, 2,$$

where (2.3) satisfies (2.2). The points i are the Bhaskara hybrid points that optimized the method, n represent the number of iterations for *-++++ step number of 2.




Figure 1: Two-step interpolation and collocation method for first order ordinary differential equations

$$
\begin{bmatrix}
1 & x_{n+1} & x_{n+1}^2 & x_{n+1}^3 & \dots & x_{n+1}^{13} & x_{n+1}^{14} \\
0 & 1 & 2x_n & 3x_n^2 & \dots & 13x_n^{12} & 14x_n^{13} \\
0 & 1 & 2x_{n+\frac{5}{26}} & 3x_{n+\frac{5}{26}}^2 & \dots & 13x_{n+\frac{5}{26}}^{12} & 14x_{n+\frac{5}{26}}^{13} \\
0 & 1 & 2x_{n+\frac{20}{29}} & 3x_{n+\frac{20}{39}}^2 & \dots & 13x_{n+\frac{20}{39}}^{12} & 14x_{n+\frac{20}{39}}^{13} \\
0 & 1 & 2x_{n+1} & 3x_{n+1}^2 & \dots & 13x_{n+1}^{12} & 14x_{n+1}^{13} \\
0 & 1 & 2x_{n+\frac{47}{26}} & 3x_{n+\frac{47}{26}}^2 & \dots & 13x_{n+\frac{47}{26}}^{12} & 14x_{n+\frac{47}{26}}^{13} \\
0 & 1 & 2x_{n+2} & 3x_{n+2}^2 & \dots & 13x_{n+2}^{12} & 14x_{n+2}^{13} \\
0 & 0 & 2 & 6x_n & \dots & 156x_n^{11} & 182x_n^{12} \\
0 & 0 & 2 & 6x_{n+\frac{20}{29}} & \dots & 156x_{n+\frac{20}{29}}^{11} & 182x_{n+\frac{20}{29}}^{12} \\
0 & 0 & 2 & 6x_{n+1} & \dots & 156x_{n+1}^{11} & 182x_{n+1}^{12} \\
0 & 0 & 2 & 6x_{n+\frac{38}{29}} & \dots & 156x_{n+\frac{38}{29}}^{11} & 182x_{n+\frac{38}{29}}^{12} \\
0 & 0 & 2 & 6x_{n+\frac{47}{26}} & \dots & 156x_{n+\frac{47}{26}}^{11} & 182x_{n+\frac{47}{26}}^{12} \\
0 & 0 & 2 & 6x_{n+2} & \dots & 156x_{n+2}^{11} & 182x_{n+2}^{12}
\end{bmatrix}
\begin{bmatrix}
\alpha_1 \\ \alpha_2 \\ \alpha_3 \\ \alpha_4 \\ \alpha_5 \\ \alpha_6 \\ \alpha_7 \\ \alpha_8 \\ \alpha_9 \\ \alpha_{10} \\ \alpha_{11} \\ \alpha_{12} \\ \alpha_{13} \\ \alpha_{14} \\ \alpha_{15}
\end{bmatrix}
=
\begin{bmatrix}
y_{n+1} \\ f_n \\ f_{n+\frac{5}{26}} \\ f_{n+\frac{20}{29}} \\ f_{n+1} \\ f_{n+\frac{38}{29}} \\ f_{n+\frac{47}{26}} \\ f_{n+2} \\ g_n \\ g_{n+\frac{5}{26}} \\ g_{n+\frac{20}{29}} \\ g_{n+1} \\ g_{n+\frac{38}{29}} \\ g_{n+\frac{47}{26}} \\ g_{n+2}
\end{bmatrix}
\tag{2.5}
$$

The system (6) was solved using the power series method, enhanced by Maple, to obtain the unknown coefficient α_j which are substituted into (1.1) to have the continuous form of the collocation method:

$$
\begin{aligned}
y(x) &= \alpha_0(x)y_{n+1} + h\sum_{j=0}^{k}\gamma_j(x)f_n + h^2\sum_{j=0}^{k}\beta_j(x)g_n \\
&+ \gamma_{\frac{5}{26}}(x)f_{n+\frac{5}{26}} + \gamma_{\frac{20}{29}}(x)f_{n+\frac{20}{29}} + \gamma_{\frac{38}{29}}(x)f_{n+\frac{38}{29}} + \gamma_{\frac{47}{26}}(x)f_{n+\frac{47}{26}} \\
&+ \beta_{\frac{5}{26}}(x)g_{n+\frac{5}{26}} + \beta_{\frac{20}{29}}(x)g_{n+\frac{20}{29}} + \beta_{\frac{38}{29}}(x)g_{n+\frac{38}{29}} + \beta_{\frac{47}{26}}(x)g_{n+\frac{47}{26}}
\end{aligned}
\tag{2.6}
$$

where $h = x_{n+1} - x_n$ and whose derivative is given by

$$
f(x) = \frac{dy}{dx}, \quad g(x) = \frac{d^2y}{dx^2} \tag{2.7}
$$

where $\alpha_j(x)$, $\gamma_j(x)$ and $\beta_j(x)$ are continuous coefficients that are uniquely determined. The main methods are gotten by evaluating (2.6) at 0, 5/26, 20/29, 38/29, 47/26, 2 to give the following:

$$
\begin{aligned}
(2.8)\qquad y_{n+\frac{5}{26}} = y_{n+1} &- \frac{13408053437210166672657 3\, h^2 g_n}{99064732150286353285120000} \\
&+ \frac{797053372151911\, h^2 g_{n+1}}{7623023165282304} \\
&+ \frac{34190738020242271746243\, h^2 g_{n+2}}{99064732150286353285120000} \\
&- \frac{1987403759042034403\, h^2 g_{n+\frac{5}{26}}}{105383133144575000000} \\
&+ \frac{16634750722379102220287979813 97\, h^2 g_{n+\frac{20}{29}}}{19918586701524685003315200000000} \\
&+ \frac{15744736418845351708654288530 07\, h^2 g_{n+\frac{38}{29}}}{59755760104574055009945600000000} \\
&+ \frac{257675377558397869\, h^2 g_{n+\frac{47}{26}}}{105383133144575000000} \\
&- \frac{99425197466903954482271423 1\, h f_n}{34024925311617582109081600000} \\
&- \frac{7437477576455479\, h f_{n+1}}{23583727917592128} \\
&- \frac{11954400821927194452844950 9\, h f_{n+2}}{17692961162041142696722432000} \\
&- \frac{11638884077093560853641 81\, h f_{n+\frac{5}{26}}}{66913223050019328125000 00} \\
&- \frac{31833672916846063655073421809594864101\, h f_{n+\frac{20}{29}}}{35891550359811098966036200320000000000} \\
&- \frac{68678001114966269287893215421630368723\, h f_{n+\frac{38}{29}}}{35891550359811098966036200320000000000} \\
&- \frac{15838025909754656597563\, h f_{n+\frac{47}{26}}}{66913223050019328125000 00}
\end{aligned}
$$

$$
\begin{aligned}
(2.9)\qquad y_{n+\frac{20}{29}} = y_{n+1} &- \frac{769431646345766277383001\, h^2 g_n}{33066146958818461463261440000} \\
&+ \frac{201236166015324651\, h^2 g_{n+1}}{14141652940152956414} \\
&+ \frac{55133896486089316940279 1\, h^2 g_{n+2}}{33066146958818461463261440000} \\
&- \frac{99718889493963361099196159 2512\, h^2 g_{n+\frac{5}{26}}}{52994555596996953229630559269531 25} \\
&- \frac{1482810138576009927\, h^2 g_{n+\frac{20}{29}}}{210959219336800000000}
\end{aligned}
$$

$$
\begin{aligned}
&+ \frac{2578074557010805647\,h^2 g_{n+\frac{38}{29}}}{1476714535357600000000}\\
&+ \frac{65761247859071138817361983 4176\,h^2 g_{n+\frac{47}{26}}}{52994555596996953229630559269 53125}\\
&- \frac{13838351177140292605769179 7697\,hf_n}{29528069234224886086692465920 0000}\\
&- \frac{414775413207763584\,hf_{n+1}}{2682037626580733113}\\
&- \frac{38985859836359753839409735 91\,hf_{n+2}}{11811227693689954434676986368000}\\
&- \frac{20849926617627333885543535033008682368\,hf_{n+\frac{5}{26}}}{61241170879834641614076444673861181640625}\\
&- \frac{95169775198571597583233 97\,hf_{n+\frac{20}{29}}}{67967059519605400000000000}\\
&- \frac{13958154599823135973713 3\,hf_{n+\frac{38}{29}}}{97095799313722000000000000}\\
&- \frac{96300581117986803401844306512677136 64\,hf_{n+\frac{47}{26}}}{61241170879834641614076444673861181640625}
\end{aligned}
$$

$$
\begin{aligned}
y_{n+1} = y_n &+ \frac{31390770194016413\,hf_n}{300800314636800000}\\
&+ \frac{24459243465753795421107944216 32\,hf_{n+\frac{5}{26}}}{85955971588987368086705566406 25}\\
&+ \frac{63191782568557371257244553881 7\,hf_{n+\frac{20}{29}}}{11209804782876811049400000000000}\\
&+ \frac{290009389184\,hf_{n+1}}{884260959813} + \frac{26934211400636986059787365139 9\,hf_{n+\frac{38}{29}}}{12455338647640901160000000000000}\\
&+ \frac{22990550579431312246763678336\,hf_{n+\frac{47}{26}}}{85955971588987368086705566406 25}\\
&+ \frac{659138643026173\,hf_{n+2}}{84224088098304000}\\
&+ \frac{706332168403\,h^2 g_n}{235789720320000}\\
&+ \frac{968545076853861104832\,h^2 g_{n+\frac{5}{26}}}{82645899987127911328125}\\
&- \frac{25436944733375565544933 3\,h^2 g_{n+\frac{20}{29}}}{27994767516705528000000 00}
\end{aligned}
\tag{2.10}
$$

$$
\begin{aligned}
&- \frac{6220321673\,h^2 g_{n+1}}{53591573322} \\
&- \frac{83896869075739821207541\,h^2 g_{n+\frac{38}{29}}}{2799476751670552800000000} \\
&- \frac{700266394943388764608\,h^2 g_{n+\frac{47}{26}}}{24793769996138373398437 5} \\
&- \frac{94340553773\,h^2 g_{n+2}}{235789720320000}
\end{aligned}
$$

$$
\begin{aligned}
(2.11)\qquad y_{n+\frac{38}{29}} = y_{n+1} &+ \frac{5513389648608931694027 91\,h^2 g_n}{330661469588184614632614 40000} \\
&+ \frac{201236166015324651\,h^2 g_{n+1}}{14141652940152956414} \\
&- \frac{76943164634576627738300 1\,h^2 g_{n+2}}{330661469588184614632614 40000} \\
&+ \frac{65761247859071138817361 9834176\,h^2 g_{n+\frac{5}{26}}}{52994555596996953229630 55926953125} \\
&+ \frac{257807455701080564 7\,h^2 g_{n+\frac{20}{29}}}{1476714535357600000000} \\
&- \frac{1482810138576009927\,h^2 g_{n+\frac{38}{29}}}{210959219336800000000} \\
&- \frac{9971888949396336109919615925 12\,h^2 g_{n+\frac{47}{26}}}{52994555596996953229630 55926953125} \\
&+ \frac{3898585983635975383940973591\,h f_n}{1181122769368995443467698 6368000} \\
&+ \frac{414775413207763584\,h f_{n+1}}{2682037626580733113} \\
&+ \frac{1383835117714029260576917976 97\,h f_{n+2}}{2952806923422488608669246592 00000} \\
&+ \frac{9630058111798680340184430651267713664\,h f_{n+\frac{5}{26}}}{6124117087983464161407644467386118164 0625} \\
&+ \frac{1395815459982313597371 33\,h f_{n+\frac{20}{29}}}{9709579931372200000000000} \\
&+ \frac{9516977519857159758323397\,h f_{n+\frac{38}{29}}}{6796705951960540000000000 0} \\
&+ \frac{2084992661762733388554353503300868236 8\,h f_{n+\frac{47}{26}}}{6124117087983464161407644467386118164 0625}
\end{aligned}
$$

$$
\begin{aligned}
(2.12)\qquad y_{n+\frac{47}{26}} = y_{n+1} &+ \frac{34190738020242271746243\,h^2 g_n}{99064732150286353285120000} \\
&+ \frac{797053372151911\,h^2 g_{n+1}}{7623023165282304} \\
&- \frac{134080534372101666726573\,h^2 g_{n+2}}{99064732150286353285120000} \\
&+ \frac{257675377558397869\,h^2 g_{n+\frac{5}{26}}}{105383133144575000000} \\
&+ \frac{157447364188453517086542885300 7\,h^2 g_{n+\frac{20}{29}}}{59755760104574055009945600000000} \\
&+ \frac{16634750722379102220287979813 97\,h^2 g_{n+\frac{38}{29}}}{19918586701524685003315200000000} \\
&- \frac{1987403759042034403\,h^2 g_{n+\frac{47}{26}}}{105383133144575000000} \\
&+ \frac{11954400821927194452844950 9\,hf_n}{17692961162041142696722432000} \\
&+ \frac{7437477576455479\,hf_{n+1}}{23583727917592128} \\
&+ \frac{99425197466903954482271423 1\,hf_{n+2}}{34024925311617582109081600000} \\
&+ \frac{15838025909754656597563\,hf_{n+\frac{5}{26}}}{66913223050019328125000000} \\
&+ \frac{68678001114966269287893215421630368723\,hf_{n+\frac{20}{29}}}{35891550359811098966036200320000000000} \\
&+ \frac{31833672916846063655073421809594864101\,hf_{n+\frac{38}{29}}}{35891550359811098966036200320000000000} \\
&+ \frac{11638884077093560853641 81\,hf_{n+\frac{47}{26}}}{66913223050019328125000000}
\end{aligned}
$$

$$
\begin{aligned}
(2.13)\qquad y_{n+2} = y_{n+1} &+ \frac{94340553773\,h^2 g_n}{235789720320000} \\
&+ \frac{6220321673\,h^2 g_{n+1}}{53591573322} \\
&- \frac{706332168403\,h^2 g_{n+2}}{235789720320000} \\
&+ \frac{700266394943388764608\,h^2 g_{n+\frac{5}{26}}}{24793769996138373398 4375} \\
&+ \frac{83896869075739821207541\,h^2 g_{n+\frac{20}{29}}}{27994767516705528000000 00} \\
&+ \frac{25436944733337556554493 33\,h^2 g_{n+\frac{38}{29}}}{27994767516705528000000 00} \\
&- \frac{968454076853861104832\,h^2 g_{n+\frac{47}{26}}}{82645899987127911328125} \\
&+ \frac{659138643026173\,h f_n}{84224088098304000} \\
&+ \frac{290009389184\,h f_{n+1}}{884260959813} \\
&+ \frac{31390770194016413\,h f_{n+2}}{300800314636800000} \\
&+ \frac{22990550579431312246763678336\,h f_{n+\frac{5}{26}}}{85955971588987368086705566406 25} \\
&+ \frac{26934211400636986059787365139 9\,h f_{n+\frac{20}{29}}}{12455338647640901166000000000 0} \\
&+ \frac{63191782568557371257244553881 7\,h f_{n+\frac{38}{29}}}{11209804782876811049400000000 00} \\
&+ \frac{24459243465753795421107944216 32\,h f_{n+\frac{47}{26}}}{85955971588987368086705566406 25}
\end{aligned}
$$

3. Analysis of the method

3.1. **Local truncation error.** The above two-step method is presented in this section. The linear operator is considered as:

$$
(3.1)\qquad L[y(x_n), h] = \sum_{i=0}^{k} \alpha_i \cdot y(x_n + ih) - h\gamma_i f(x_n + ih) + h^2 \beta_i g(x_n + ih)
$$

The function $y(x)$ is an arbitrary test function that is continuously differentiable in the interval $[a, b]$. Expanding $y(x_n + ih)$, $f(x_n + ih)$ and $g(x_n + ih)$ in Taylors series about x_n and factoring the coefficients of h to get

$$L[y(x_n), h] = C_0 y(x_n) + C_1 h y(x_n) + C_2 h^2 y^2(x_n) + \ldots + C_p h^p y^p(x_n) + \ldots \tag{3.2}$$

where the constant C_i, $i = 0, 1, \ldots$ are given as follows

$$\begin{cases} C_0 &= \alpha_0 + \alpha_1, \\ C_1 &= \alpha_1 - (\gamma_0 + \gamma_{\frac{5}{26}} + \gamma_{\frac{20}{29}} + \gamma_1 + \gamma_{\frac{38}{29}} + \gamma_{\frac{47}{26}} + \gamma_2), \\ C_2 &= \frac{1}{2!}\alpha_i - \left(\frac{5}{26}\gamma_{\frac{5}{26}} + \frac{20}{29}\gamma_{\frac{20}{29}} + \gamma_1 + \frac{38}{29}\gamma_{\frac{38}{29}} + \frac{47}{26}\gamma_{\frac{47}{26}} + 2\gamma_2\right) \\ &\quad - \left(\beta_0 + \beta_{\frac{5}{26}} + \beta_{\frac{20}{29}} + \beta_1 + \beta_{\frac{38}{29}} + \beta_{\frac{47}{26}} + \beta_2\right) \\ \ldots & \\ C_p &= \frac{1}{p!}\alpha_1 - \frac{1}{(p-1)!}\left(\left(\frac{5}{26}\right)^{p-1}\gamma_{\frac{5}{26}} + \left(\frac{20}{29}\right)^{p-1}\gamma_{\frac{20}{29}} + \gamma_1 + \left(\frac{38}{29}\right)^{p-1}\gamma_{\frac{38}{29}}\right. \\ &\quad \left. + \left(\frac{47}{26}\right)^{p-1}\gamma_{\frac{47}{26}} + 2^{p-1}\gamma_2\right) \\ &\quad - \frac{1}{(p-2)!}\left(\left(\frac{5}{26}\right)^{p-2}\beta_{\frac{5}{26}} + \left(\frac{20}{29}\right)^{p-2}\beta_{\frac{20}{29}} + \gamma_1 + \left(\frac{38}{29}\right)^{p-2}\beta_{\frac{38}{29}}\right. \\ &\quad \left. + \left(\frac{47}{26}\right)^{p-2}\beta_{\frac{47}{26}} + 2^{p-2}\beta_2\right) \end{cases} \tag{3.3}$$

From (3.2), we can obtain $C_0 = C_1 = \ldots = C_{14} = 0$ and

$$\begin{aligned} C_{15} \;=\; \Bigg(&\frac{742953457}{254252803376756856672000}, \\ &\frac{1898244635071516210773}{776272612894431197030421294481408000}, \\ &\frac{373071477459644081757}{42551629774433552891797727262934720000}, \\ &\frac{1898244635071516210773}{776272612894431197030421294481408000}, \\ &\frac{742953457}{254252803376756856672000}\Bigg)^{\top}. \end{aligned}$$

Therefore the proposed method is of order 14 [8].

3.2. **Stability Analysis.** (2.8) can be written in the matrix form

$$A_1 Y_{n+1} = A_0 Y_n + h(B_0 F_n + B_1 F_{n+1}) + h^2(C_0 G_n + C_1 G_{n+1}) \tag{3.4}$$

where A_0, A_1, B_0, B_1, C_0, C_1 are 6×6 matrix and

$Y_n = (y_n, y_{n-\frac{5}{26}}, y_{n-\frac{20}{29}}, y_{n-\frac{38}{29}}, y_{n-\frac{47}{26}}, y_{n-2})^{\top}$,

$Y_{n+1} = (y_{n+\frac{5}{26}}, y_{n+\frac{20}{29}}, y_{n+1}, y_{n+\frac{38}{29}}, y_{n+\frac{47}{26}}, y_{n+2})^{\top}$,

$F_n = (f_n, f_{n-\frac{5}{26}}, f_{n-\frac{20}{29}}, f_{n-\frac{38}{29}}, f_{n-\frac{47}{26}}, f_{n-2})^{\top}$,

$F_{n+1} = (f_{n+\frac{5}{26}}, f_{n+\frac{20}{29}}, f_{n+1}, f_{n+\frac{38}{29}}, f_{n+\frac{47}{26}}, f_{n+2})^{\top}$,

$G_n = (g_n, g_{n-\frac{5}{26}}, g_{n-\frac{20}{29}}, g_{n-\frac{38}{29}}, g_{n-\frac{47}{26}}, g_{n-2})^{\top}$,

$G_{n+1} = (g_{n+\frac{5}{26}}, g_{n+\frac{20}{29}}, g_{n+1}, g_{n+\frac{38}{29}}, g_{n+\frac{47}{26}}, g_{n+2})^{\top}$.

According to Lambert [11], a numerical technique is considered to be zero-stable if no root of the first characteristics polynomial $\rho(z)$ has a modulus more than one, and every root of modulus one has simplicity not greater than the order of the differential equation.

The characteristics equation of the hybrid block method in (3.3) is

$$\rho(z) = det(ZA_1 - A_0) = z^5(z+1) = 0. \tag{3.5}$$

This implies that $z_1 = z_2 = z_3 = z_4 = z_5 = 0$ and $|z_6| = 1$. Therefore, the method is zero stable.

A method is consistency if it has order greater than or equal to 1. Therefore, the proposed method is consistency Jator [9].

In addition, the method is converging only if it is consistent and zero stable Henrici [8]. Because the proposed method meets the two requirements, it is convergent.

The absolute stability region was studied. The linear stability properties of the proposed method are studied by applying it to the test problem $y' = \lambda y$ with $\lambda < 0$ to get

$$Y_{n+1} = M(z)Y_n, \quad z = \lambda h \tag{3.6}$$

where

$$M(z) = (A_1 - zB_1 - z^2C_1)^{-1}(A_0 + zB_0 + z^2C_0). \tag{3.7}$$

The matrix $\sigma(z)$ has eigenvalues $\{0, 0, 0, ..., \lambda_k\}$, and the dominant eigenvalue $\lambda_k : \mathrm{C} \to \mathrm{C}$ is a rational function (called the stability function) with real coefficients given by

$$\lambda_k = \frac{P(z)}{P(-z)}. \tag{3.8}$$

It is clear from the stability functions that for $Re(z) < 0$, $|\lambda_k| \leq 1$.

The absolute stability region is given Figure 2. The stability region contains the entire left half complex plane and thus, the method is A-stable [2].




Figure 2: Absolute Stability Region of the Method

3.3. **Convergence of the method.** Consistency and zero stability are both required and sufficient for a linear multistep technique to reach convergence in the spirit of Lambert [11]. As a result, we infer that our technique is convergent since it has an order of accuracy (which implies consistency) and zero–stable.

4. Numerical experiments

The efficiency of the method will be demonstrated using some examples.

Example 1. Consider the following nonlinear equation:

$$y' + y = 2t + e^t, \quad y(0) = -1$$

with exact solution

$$y(t) = \cosh(t) + 2t - 2.$$

Table 1 shows the comparison between the proposed method and Motsa [16]. Comparison between the numerical solutions and the exact solutions is shown in Figure 3 when $h = 0.1$.

Table 1: Comparison between the absolute errors in the proposed method and the method in paper [16].

t	Error in our method	Error in [16], $M = 5$
0.0	0.000	0.0000
0.1	2.62E-31	1.40E-15
0.2	5.33E-31	5.44E-15
0.3	1.30E-30	1.19E-14
0.4	2.05E-30	2.07E-14
0.5	3.27E-30	3.17E-14
0.6	4.44E-30	4.47E-14
0.7	6.08E-30	5.97E-14
0.8	7.66E-30	7.69E-14
0.9	9.72E-30	9.61E-14
1.0	1.17E-29	1.17E-13
1.1	1.42E-29	1.41E-13
1.2	1.67E-29	1.67E-13
1.3	1.97E-29	1.96E-13
1.4	2.27E-29	2.27E-13
1.5	2.63E-29	2.61E-13
1.6	2.98E-29	2.99E-13
1.7	3.42E-29	3.40E-13
1.8	3.85E-29	3.85E-13
1.9	4.37E-29	4.34E-13
2.0	4.88E-29	4.88E-13




Figure 3: Exact solution and Numerical solution for Example 1

Example 2. Consider the following nonlinear equation

$$y' = y^2 - ty, \quad y(0) = \frac{1}{2}$$

with exact solution

$$y(t) = \frac{2\exp\{-t^2\}}{\sqrt{\pi}t^2 - 4}.$$

The exact and numerical solutions for Example 2 are displayed in Figure 4. The errors comparison with other methods are illustrated in Table 2.

Table 2: Numerical, exact solutions, and the absolute errors.

t	Solution by our method	Exact solution	Absolute Error
0.0	0.50000	0.50000	0.00000
1.0	0.530010126	0.530010126	2.04E-22
2.0	0.168387788	0.168387788	5.36E-22
3.0	1.48E-02	1.48E-02	1.40E-23
4.0	4.49E-04	4.49E-04	6.73E-26
5.0	4.99E-06	4.99E-06	1.05E-26




Figure 4: Exact solution and Numerical solution for Example 2

Example 3. Consider the Riccati equation. A matrix equation with an analogous quadratic term is referred to as Riccati equation, this matrix occurs in both discrete-time and continuous-time linear-quadratic-gaussian control.

$$y' = -y^2 + 2y + 1, \quad y(0) = 0,$$

with exact solution

$$y(t) = 1 + \sqrt{2}\tanh\sqrt{2}t + \frac{1}{2}\log\frac{\sqrt{2}-1}{\sqrt{2}+1}.$$

The exact and numerical solutions are presented in the Table 3.

Table 3: Comparison between the absolute errors in our method and the methods in [10, 16].

t	Error in our method	Error in [16], $M = 5$	Error in [10]
0	0.0000	0.0000	0.0000
1	1.79E-23	3.54E-13	1.61E-10
2	2.21E-24	3.07E-14	2.14E-10
3	3.96E-24	8.08E-15	2.22E-10
4	3.73E-24	1.74E-15	2.14E-10
5	4.00E-24	1.81E-16	1.99E-10
6	4.72E-24	1.53E-17	1.82E-10
7	4.37E-24	1.18E-18	1.65E-10
8	3.96E-24	8.60E-20	1.50E-10
9	3.50E-24	6.04E-21	1.37E-10
10	3.75E-24	4.14E-22	1.25E-10




Figure 5: Exact solution and Numerical solution for Example 3

Example 4. Consider the Lorenz system equation are derived to model some unpredictable behavior of weather.

$$\begin{cases} y_1' = a(y_2 - y_1), & y_1(0) = 1, \\ y_2' = -y_1 y_3 + b y_1 - y_2, & y_2(0) = 5, \\ y_3' = y_1 y_3 - c y_3, & y_3(0) = 10. \end{cases}$$

The constants are $a = 10$, $b = 28$, and $c = 8/3$. The phase portraits for the Lorenz equation are displayed in Figure 6.




Figure 6: Phase portraits for the Lorenz system, $h = 0.01$

Example 5. Consider the chaotic the chaotic system called the Arnodo-Coullet which is system given by the following set of equations

$$\begin{cases} y_1' = y_2, & y_1(0) = 0.21, \\ y_2' = y_3, & y_2(0) = 0.22, \\ y_3' = ay_1 - by_2 - y_3 - y_1^3, & y_3(0) = 0.61. \end{cases}$$

The constants are $a = 5$, $b = 3.8$. The phase portraits for the Arnodo-Coullet are displayed in Figure 7.




Figure 7: Phase portraits for the Arnodo-Coullet system, $h = 0.025$

Discussion of Results. Example 1 is a non-linear equation, the result from our method was compared to paper [16], our method performs better than that of [16] which has 4.88×10^{-29} as error compared to 1.40×10^{-15} of [16]. Example 2 is a non-linear equation, we compared the result with the exact solution which compare favorably well with the exact solution. Example 3 is a Riccati equation (this matrix occurs in both discrete-time and continuous-time linear-quadratic-gaussian control) as shown in Table 3, our method performs significantly better than papers [16] and [10]. The Lorenz system equation of Example 4 are derived to model some unpredictable behavior of weather, The phase portraits shows the graphical solution of the problem that arises from chaos theory. Lastly, we considered the chaotic system called the Arnodo-Coullet which is system given by a set of equations, The phase portraits also demonstrates the solution of the problem solved by our method. It is clearly demonstrated from the examples that the proposed method is able to provide solution to non-linear dynamical problems.

Conclusion. For the self-starting solution of general first initial value problems of ordinary differential equations, we provide an optimized two-step block technique using Bhaskara hybrid points for non-linear dynamical first order differential equations. The method's performance was assessed using four Bhaskara hybrid points. For orders up to 14, the suggested hybrid block technique is both zero stable and converging. Riccati equation, chaotic system dubbed Arnodo-Coullet and non-linear stiff issue are all solved by the suggested approach. The absolute inaccuracy falls as h decreases in these situations. The numerical examples demonstrate the method's effectiveness and accuracy in comparison to other approaches in the literature.

Acknowledgement: The authors are grateful to Federal University of Technology, Minna, Nigeria, National Open University of Nigeria and Tomsk Polytechnics University, Russia for the supports they received during the compilation of this work.

Competing interests: The manuscript was read and approved by all the authors. They therefore declare that there is no conflicts of interest.

Funding: The Authors received no financial support for the research, authorship, and/or publication of this article.

References

[1] Akinfenwa O. A., Abdulganiy R. I., Okunuga S. A. & Irechukwu V. (2017). Simpson's 3/8 type block method for stiff systems of ordinary differential equations. *Journal of the Nigerian Mathematical Society.* **36** (3), 503-514.

[2] Akinfenwa O. A., Akinnukawe B. & Mudasiru S. B. (2015). A family of continuous third derivative block methods for solving stiff systems of first order ordinary differential equations. *J. Nigerian Mathematical Society.* **34** (2), 160-168.

[3] Akinfenwa O. A., Jator S. N. & Yao N. M. (2011). A Self-starting Block Adams Methods for Solving Stiff ODEs. *The 14th IEEE International Conference on Computational Science and Engineering.* **156**, 127-136.

[4] Areo E. A. & Adeniyi R. B. (2014). Block implicit one-step method for the numerical integration of initial value problems in ordinary differential equations. *Int J Math Stat Stud.* **2** (3). 4-13.

[5] Badmus A. M., Yahaya Y. A. & Subair A. O. (2014). Implicit Stomer Cowell k-step hybrid block methods for solution of first order ordinary differential equations (ODEs). *Int. J. Appl. Math. Res.* **3** (4), 464-72.

[6] Fotta A. U., Alabi T. J. (2015). Block method with one hybrid point for the solution of first order initial value problems of ordinary differential equations. *Int. J. Pure Appl. Math.* **103** (3), 511-521.

[7] Garba J. & Mohammed U. (2020). Derivation of a new one-step numerical integrator with three intra-step points for solving the first order ordinary differential equations. *Nigeria Journal of Mathematics and Applications.* **30**, 155-172.

[8] Henrici P. (1962). *Discrete variable methods in ODE.* New York, NY, USA: John Wiley & Sons.

[9] Jator S. (2015). A sixth order linear multistep method for the direct solution of $y = f(x, y, y')$. *Int. J. Pure Appl. Math.* **40** (7), 457-72.

[10] Kashkari S. H. B. & Syam M. I. (2019). Optimization of one step block method with three hybrid points for solving first-order ordinary differential equations. *Results in Physics.* **12**, 592-596.

[11] Lambert J. D. (1973). *Computational methods in ODE.* New York, NY, USA: John Wiley & Sons.

[12] Mehdizadeh K. & Molayi M. (2015). The New Class of A-Stable Hybrid Multistep Methods for Numerical polution of Stiff Initial Value Problem. *Mathematical Theory and Modeling.* **5** (1), 97-103.

[13] Milne W. E. (1953). *Numerical solution of differential equations.* New York, NY, USA: John Wiley & Sons.

[14] Mohammed U., Garba J. & Semenov M. E. (2021). One-step second derivative block intra-step method for stiff system of ordinary differential equations. *Journal of Nigerian Mathematical Society.* **40** (1), 74-77.

[15] Mehdizadeh K., Nasehi N. & Hojjati G. (2012). A Class of Second Derivative Multistep Methods for Stiff Systems. *Acta Universitatis Apulensis.* **30**, 171-188.

[16] Motsa S. S. (2021). *(Spectral) Block hybrid methods.* UK-Africa Postgraduate Study Institute in Mathematical Sciences.

[17] Ramos H. (2017). An optimized two-step hybrid block method for solving first-order initialvalue problems in ODEs derivation of the block hybrid method. *Diff. Geometry DynSyst.* **19**, 107-118.

[18] Ramos H., Kalogiratou Z., Monovasilis T. & Simos T. E. (2016). An optimized twostep hybrid block method for solving general second order initial-value problems. *Numerical Algorithms.* **72** (4), 1089-1102.

[19] Robertson1966 Robertson H. H. (1966). *The Solution Of A Set Of Reaction Rate Equations.* Academic Press.

[20] Shampine L. F. & Watts H. A. (1969). Block Implicit One-Step Methods. *Math. Comp.* **23**, 731-740.

[21] Sunday J., Skwame Y. & Tumba P. (2015). A quarter-step hybrid block method for first-order ordinary differential equations. *British J. Math. Comput. Sci.* **6** (4), 269-78.

[22] Yahaya Y. A. & Tijjani A. A. (2015). Formulation of corrector methods from 3-step hybrid Adams type methods for the solution of first order ordinary differential equation. 102-107.
[23] Yakusak N. S. & Adeniyi R. B. (2015). A four-step hybrid block method for first order initial value problems in ordinary differential equations. *AMSE J.* **52** (1), 17-30.