DOI: https://doi.org/10.67897/njma.2022.hfxgvhz7



Nigerian Journal of Mathematics and Applications
Volume **32**, (2022), 80 – 89. *Printed by Unilorin press*
©*Nig. J. Math. Appl.* *http : //www.njmaman.com*

Stability of Solutions for a Class of Second Order Nonlinear Integro-Differential Equations with Delay

Daniel Oluwasegun Adams[1*], Rasheed Ayodeji Adetona[2], Musiliu Tayo Raji[3], Samson Abidemi Oni[3] and Akinwale Lewis Olutimo[4]

Abstract

In this paper, we investigate the stability of solutions for a class of second order nonlinear delay integro-differential equations by using a suitable Lyapunov-Krasovskiĭ functional with sufficient conditions to establish some new results. An example is given to show the validity of the results obtained.

1. Introduction

In relevant literature, the mathematical model known as Volterra integro-differential equation which appeared after its establishment by Vito Volterra in 1926 is viable and its application is increasing to various fields ([25], [27]). Volterra integro-differential equations are important effective mathematical models used to describe many real world phenomena concerning atomic energy, biology, chemistry, control theory, economy, engineering technique fields, information theory, medicine, population dynamics and so on. For a survey on integro-differential equations, see the following references: Adeyanju et.al [3], Ahmad and Stamova (Eds.) [4], Burton [6], Burton and Mahfoud [7], Corduneanu [8], Corduneanu and

Received: 20/04/2022, Accepted: 29/05/2022, Revised: 25/06/2022. * Corresponding author.
2015 *Mathematics Subject Classification.* 34K20 & 37C75.
Keywords and phrases. Stability; Lyapunov-Krasovskiĭ functional; integro-differential equations of second order; Nonlinear
[1&3]Department of Mathematics, Federal University of Agriculture, Abeokuta, Nigeria
[2]Department of Mathematics, Obafemi Awolowo University, Ile-Ife, Nigeria
[4]Department of Mathematics, Lagos State University, Ojo, Nigeria
E-mail of the corresponding author: danielogic2008@yahoo.com
ORCID of the corresponding author: 0000-0002-4137-1858

Sandberg (Eds.) [9], Da Prato and Iannelli (Eds.) [11], Gripenberg et.al [14], Lakshmikantham and Mohana [15], Peschel and Mende [19], Staffans [23], Tunç and Tunç [25] and Wazwaz [26]. Importantly, second order integro-differential equations appear in stability problems of visco elastic shells (see [5], [12]).
There are various research articles that have been devoted to stability of the first and second order integro-differential equations which include but not limited to the following references: [3], [5], [10], [13], [16], [17], [18], [21], [22], [24] and [27] while there are numerous published papers which considered other aspects of differential equations using Lyapunov's direct method to establish their results, see [1], [2] and [20] to mention a few.
Furthermore, in 2018 Zhao and Meng [28] considered a kind of nonlinear delay integro-differential equations

$$x'' + f(t,x,x')x' + g(t,x,x') + h(x(t-\tau)) = p(t,x(t))\int_0^t q(s,x'(s))ds \tag{1}$$

and

$$x'' + f(t,x,x')x' + g(t,x,x') + h(x) = p(x(t-\tau))\int_0^t q(s,x'(s))ds, \tag{2}$$

where $\mathbb{R}^+ = [0,\infty)$, $\tau > 0$ is a constant while f, g, h, p and q are continuous functions with respect to the argument displayed explicitly. The authors by using different Lyapunov funtionals, established the zero solutions for each of the integro-differential equation with delay and also examined the stability of the generalized form for each of (1) and (2) with variable delay $\tau(t)$ respectively.
Motivated by the work of Zhao and Meng [28], our aim is to obtain sufficient conditions for the stability of solutions for a class of nonlinear delay integro-differential equations by using a suitable Lyapunov-Krasovskiĭ functional with sufficient criteria to achieve the new results.
In this work, we consider the nonlinear second order delay integro-differential equation of the form:

$$x'' + a(t)f(t,x,x')x' + g(t,x,x') + h(x(t-\tau)) = p(x(t-\tau))\int_0^t q(s,x'(s))ds, \tag{3}$$

where $\mathbb{R}^+ = [0,+\infty)$, $\tau > 0$ is a constant and $a : \mathbb{R}^+ \longrightarrow \mathbb{R}^+$ while $f, g : \mathbb{R}^+\times\mathbb{R}^2 \longrightarrow \mathbb{R}$ and $h : \mathbb{R}^+ \longrightarrow \mathbb{R}$ are continuous with $h(0) = 0$, $p, q : \mathbb{R}^+\times\mathbb{R} \longrightarrow \mathbb{R}$ with $p(0) = 0$ and $q(t,0) = 0$. Also, $'$ implies differentiation with respect to t. Indeed, the integro-differential equation (3) extends and improves that of [28] and it is different from those in the references. Also, we provide an example to show the validity of our results.

2. Preliminary Definitions

We give some basic definitions on the stability of solution of integro-differential equations (see [3] and [6]). Consider the system of first order non-linear and non-homogeneous Voltera integro-differential equation

$$X'(t) = A(t)X(t) + \int_{t-\tau}^{t} F(t,\phi,h(X(\phi)))g(X(\phi))d\phi + E(t,X(t)), \tag{4}$$

where $t \in [0,\infty)$, $X \in \mathbb{R}^n$, $A(t)$, $F(t,\phi,h(X(\phi)))$ and $E(t,X(t))$ are continuous functions in their respective arguments explicitly displayed, such that $0 \leq \phi \leq t < \infty$, $h(0) = 0$, $h(X) \neq 0$, $X \neq 0$, $F(t,\phi,0) = 0$; $h, g : \mathbb{R}^n \longrightarrow \mathbb{R}^n$, $g(0) = 0$ are continuous functions and $\tau > 0$ is a constant delay.
Let $X(t,t_0,B)$, $t \geq t_0$ be a solution of (4) on $[t_0-\tau,\beta)$, $\beta > 0$ such that $X(t) = B(t)$ on $B \in [t_0-\tau,t_0]$ and $\|B(t)\| = sup_{t\in[t_0-\tau,t_0]}\|B(t)\|$, where $B : [t_0-\tau,t_0] \longrightarrow \mathbb{R}^n$ is a continuous initial function.
Then, we have the following definitions:

Definition 2.1. The zero solution of (4) is said to be stable if for each $\epsilon > 0$ and $t_0 \geq 0$, there exists a $\delta = \delta(t_0,\epsilon) > 0$ such that if $\|B(t)\| < \delta$ on $[t_0-\tau,t_0]$, we have $\|X(t,B)\| < \epsilon$, for all $t \geq t_0$.

Definition 2.2. The zero solution of (4) is said to be uniformly stable if δ is independent of t_0.

Definition 2.3. The zero solution of the (4) is said to be asymptotically stable if it is stable and for each $t_0 \geq 0$, there is a $\delta > 0$ such that $t \geq t_0$, $\|B(t)\| < \delta$ on $[0,t_0]$ implies $\|X(t,B)\| \longrightarrow 0$ as $t \longrightarrow \infty$.

3. Main Results

In this section, we give our main results.
We rewrite (3) as the system

$$\begin{gathered} x' = y \\ y' = -a(t)f(t,x,y)y - g(t,x,y) - h(x) + \int_{t-\tau}^{t} h'(x(s))y(s)ds \\ -p(x)\int_0^t q(s,y(s))ds + \int_{t-\tau}^{t} p'(x(s))y(s)ds \int_0^t q(s,y(s))ds. \end{gathered} \tag{5}$$

Now, we have the following Theorem.

Theorem 3.1. *Consider the system (5) above. There exist positive constants K_1, K_2, M, N, Ω_1, Ω_2, Φ, δ, τ and functions $P(t), Q(t) : \mathbb{R}^+ \longrightarrow \mathbb{R}^+$ such that the following conditions hold:*

(c_1): *$a(t) \geq \alpha > 0$, $t \in \mathbb{R}^+$;*

(c_2): $\dfrac{h(x)}{x} \geq \delta$ *when* $x \neq 0$, *for all* $x \in \mathbb{R}$, δ *a constant;*

(c_3): $|h'(x)| \leq K_1$, $|p'(x)| \leq K_2$ *and* $|q(t,y)| \leq Q(t)|y|$ *for* $y \neq 0$, *and* $x, y \in \mathbb{R}$;

(c_4): $f(t,x,0) = 0$, $g(t,x,0) = 0$, $\dfrac{f(t,x,y)}{y^2} \geq M$, $\dfrac{g(t,x,y)}{y} \geq N$ *when* $y \neq 0$ *for all* $x, y \in \mathbb{R}$;

(c_5): $|p(x)| \leq P(t)$, $\Omega_1 \leq P(t) \leq \Omega_2$, $Q(t) \leq \Phi$, $t \in \mathbb{R}^+$;

(c_6): $\dfrac{K_2\tau}{\alpha}\displaystyle\int_0^\infty Q(s)ds \leq 2M$;

(c_7): $\Omega_2 \displaystyle\int_0^\infty Q(s)ds + \frac{(\Omega_2 + K_2\tau)}{\Omega_1}\Phi \int_0^\infty P(s)ds$
$+K_1\Big(\tau - y^{-2}\displaystyle\int_{t-\tau}^t y^2(s)ds\Big) \leq 2N$ *when* $y \neq 0$;.

Then, the zero solution of system (5) is stable.

Proof. We define the Lyapunov-Krasovskiĭ functional $V(t,x,y) = V(t)$ as follows:

$$V(t,x,y) = \frac{1}{2}y^2 + \int_0^x h(\phi)d\phi + \lambda\int_{-\tau}^0 ds\int_{t+s}^t y^4(\theta)d\theta + \mu\int_0^t ds\int_t^\infty |p(x(\theta))|Q(s)y^2(s)d\theta,$$

where λ and μ are two positive constants to be determined. So, since we have

$$\int_0^x \frac{h(\phi)}{\phi}\,\phi d\phi \geq \int_0^x \delta\,\phi d\phi = \frac{\delta}{2}x^2.$$

Then,

$$V(t,x,y) \geq \frac{\delta}{2}x^2 + \frac{1}{2}y^2 > 0$$

which shows that the functional is positive definite.
Differentiating $V(t)$ along the solution path of (5), we have

$$\begin{aligned}\frac{dV(t)}{dt} &= yy' + h(x)y + \lambda\tau y^4 - \lambda\int_{t-\tau}^t y^4(s)ds + \mu Q(t)y^2(t)\int_t^\infty |p(x(\theta))|d\theta \\ &\quad -\mu|p(x(t))|\int_0^t Q(s)y^2(s)ds \\ &= -a(t)f(t,x,y)y^2 - g(t,x,y)y + \lambda\tau y^4 + y\int_{t-\tau}^t h'(x(s))y(s)ds \\ &\quad -p(x)y\int_0^t q(s,y(s))ds + y\int_{t-\tau}^t p'(x(s))y(s)ds\int_0^t q(s,y(s))ds \\ &\quad -\lambda\int_{t-\tau}^t y^4(s)ds + \mu Q(t)y^2\int_t^\infty |p(x(\theta))|d\theta - \mu|p(x(t))|\int_0^t Q(s)y^2ds. \qquad (6)\end{aligned}$$

Applying conditions (c_3) and the inequality $2ab \leq a^2 + b^2$, we have the following:

$$y(t)\int_{t-\tau}^{t} h'(x(s))y(s)ds \leq |y(t)|\int_{t-\tau}^{t} |h'(x(s))||y(s)|ds$$

$$\leq \frac{K_1}{2}\int_{t-\tau}^{t}[y^2(t)+y^2(s)]ds$$

$$= \frac{K_1}{2}\tau y^2 + \frac{K_1}{2}\int_{t-\tau}^{t} y^2(s)ds. \tag{7}$$

$$y\int_{t-\tau}^{t} p'(x(s))y(s)ds\int_{0}^{t} q(s,y(s))ds \leq |y(t)|\int_{t-\tau}^{t} |p'(x(s))||y(s)|ds\int_{0}^{t}|q(s,y(s))|ds$$

$$\leq \frac{K_2}{2}\int_{t-\tau}^{t}[y^2(t)+y^2(s)]ds\int_{0}^{t} Q(\theta)|y(\theta)|d\theta$$

$$\leq \frac{K_2}{4}\tau y^4\int_{0}^{\infty} Q(\theta)d\theta + \frac{K_2}{2}\tau\int_{0}^{t} Q(s)y^2(s)ds$$

$$+ \frac{K_2}{4}\int_{t-\tau}^{t} y^4(s)ds\int_{0}^{\infty} Q(s)ds. \tag{8}$$

$$-yp(x(t))\int_{0}^{t} q(s,y(s))ds \leq |y(t)||p(x(t))|\int_{0}^{t}|q(s,y(s))|ds$$

$$\leq |y(t)||p(x(t))|\int_{0}^{t} Q(s)|y(s))|ds$$

$$\leq \frac{1}{2}|p(x(t))|\int_{0}^{t} Q(s)[y^2(t)+y^2(s)]ds$$

$$\leq \frac{|p(x(t))|}{2}y^2\int_{0}^{\infty} Q(s)ds + \frac{|p(x(t))|}{2}\int_{0}^{t} Q(s)y^2(s)ds. \tag{9}$$

Substituting inequalities (7), (8) and (9) in equation (6), we now have

$$V'(t) \leq -a(t)f(t,x,y)y^2 - g(t,x,y)y + \lambda\tau y^4 + \frac{K_1}{2}\tau y^2 + \frac{K_1}{2}\int_{t-\tau}^{t} y^2(s)ds$$

$$+\frac{|p(x(t))|}{2}y^2\int_{0}^{\infty} Q(s)ds + \frac{|p(x(t))|}{2}\int_{0}^{t} Q(s)y^2(s)ds + \frac{K_2}{4}\tau y^4\int_{0}^{\infty} Q(\theta)d\theta$$

$$+\frac{K_2}{2}\tau\int_{0}^{t} Q(s)y^2(s)ds + \frac{K_2}{4}\int_{t-\tau}^{t} y^4(s)ds\int_{0}^{\infty} Q(s)ds - \lambda\int_{t-\tau}^{t} y^4(s)ds$$

$$+\mu Q(t)y^2\int_{t}^{\infty}|p(x(\theta))|d\theta - \mu|p(x(t))|\int_{0}^{t} Q(s)y^2(s)ds.$$

Using the conditions (c_1), (c_4) and (c_5), we obtain

$$V'(t) \leq -\Big[\alpha M - \frac{K_2}{4}\tau \int_0^\infty Q(s)ds - \lambda\tau\Big]y^4 - \Big[N - \frac{\Omega_2}{2}\int_0^\infty Q(s)ds - \mu\Phi\int_0^\infty P(s)ds - \frac{K_1}{2}\tau\Big]y^2$$
$$+\Big[\frac{K_2}{4}\int_0^\infty Q(s)ds - \lambda\Big]\int_{t-\tau}^t y^4(s)ds + \Big[\frac{\Omega_2}{2} + \frac{K_2}{2}\tau - \mu\Omega_1\Big]\int_0^t Q(s)y^2(s)ds$$
$$+\frac{K_1}{2}\int_{t-\tau}^t y^2(s)ds.$$

Hence, we choose $\lambda = \dfrac{K_2}{4}\displaystyle\int_0^\infty Q(s)ds$, $\mu = \dfrac{(\Omega_2 + K_2\tau)}{2\Omega_1}$ and by (c_6), c_7, it is clear that

$$V'(t) \leq -\Big[\alpha M - \frac{K_2}{2}\tau\int_0^\infty Q(s)ds\Big]y^4 - \Big[N - \frac{\Omega_2}{2}\int_0^\infty Q(s)ds$$
$$-\frac{(\Omega_2 + K_2\tau)}{2\Omega_1}\Phi\int_0^\infty P(s)ds - \frac{K_1}{2}\tau\Big]y^2$$
$$+\frac{K_1}{2}\int_{t-\tau}^t y^2(s)ds \leq 0.$$

Thus, the zero solution of system (5) is stable and the proof completed.

Remark. If $h(x(t-\tau)) = h(x)$, where $t > \tau$, $\tau > 0$ for $t \in \mathbb{R}^+$ and $x \in \mathbb{R}$. Then equation (3) reduces to the one considered for Theorem 3 in Zhao and Meng [28].

We now generalise equation (3) to the form with a variable delay $\tau(t)$. Such that its equivalent system now become

$$x' = y,$$
$$y' = -a(t)f(t,x,y)y - g(t,x,y) - h(x) + \int_{t-\tau(t)}^t h'(x(s))y(s)ds$$
$$-p(x)\int_0^t q(s,y(s))ds + \int_{t-\tau(t)}^t p'(x(s))y(s)ds\int_0^t q(s,y(s))ds. \tag{10}$$

We now give some new conditions to obtain additional result.

Corollary. Let conditions (c_1) to (c_5) of the above Theorem hold. Consider the system (10) with a variable delay $\tau(t)$ satisfying the following additional conditions:

(c_8)**:** there are $\tau > 0$ and $0 < \beta < 1$, such that $0 \leq \tau(t) \leq \tau$ and $\tau'(t) \leq \beta$;

(c_9)**:** $\dfrac{K_2\tau}{\alpha}\cdot\dfrac{(2-\beta)}{(1-\beta)}\displaystyle\int_0^\infty Q(s)ds \leq 4M.$

Then, the zero solution of system (10) with a variable delay $\tau(t)$ is stable.

Proof. Though the proof is similar to that of the above Theorem. But now, with the application of condition (c_8) we have

$$V'(t) \leq -\Big[\alpha M - \frac{K_2}{4}\tau \int_0^\infty Q(s)ds - \lambda\tau\Big]y^4 - \Big[N - \frac{\Omega_2}{2}\int_0^\infty Q(s)ds - \mu\Phi\int_0^\infty P(s)ds - \frac{K_1}{2}\tau\Big]y^2$$
$$+\Big[\frac{K_2}{4}\int_0^\infty Q(s)ds - \lambda(1-\beta)\Big]\int_{t-\tau}^t y^4(s)ds + \Big[\frac{\Omega_2}{2} + \frac{K_2}{2}\tau - \mu\Omega_1\Big]\int_0^t Q(s)y^2(s)ds$$
$$+\frac{K_1}{2}\int_{t-\tau}^t y^2(s)ds.$$

Now, we choose $\lambda = \dfrac{K_2}{4(1-\beta)}\displaystyle\int_0^\infty Q(s)ds$, $\mu = \dfrac{(\Omega_2 + K_2\tau)}{2\Omega_1}$ and by (c_7), (c_9), it is obvious that

$$V'(t) \leq -\Big[\alpha M - \frac{(2-\beta)}{4(1-\beta)}K_2\tau\int_0^\infty Q(s)ds\Big]y^4 - \Big[N - \frac{\Omega_2}{2}\int_0^\infty Q(s)ds$$
$$-\frac{(\Omega_2 + K_2\tau)}{2\Omega_1}\Phi\int_0^\infty P(s)ds - \frac{K_1}{2}\tau\Big]y^2$$
$$+\frac{K_1}{2}\int_{t-\tau}^t y^2(s)ds \leq 0.$$

3.1. **Example.** To show the effectiveness of the results, we consider the following second order nonlinear delay integro-differential equation of the form:

$$x''(t) + 5(t + x^2(t) + x'^2(t))x'(t) + \frac{1}{4}x'(t) + 5x(t-1) = e^{-2(t-1)}\int_0^t e^{-s}x'(s)ds \tag{11}$$

which is a special case of equation (3). This equation (11) can be re-written as follows:
Let $x'(t) = y(t)$, then

$$y'(t) + 5(t + x^2(t) + y^2(t))y(t) + \frac{1}{4}y(t) + 5x(t-1) - e^{-2(t-1)}\int_0^t e^{-s}y(s)ds = 0. \tag{12}$$

In the example above,
$f(t,x,y) = (t + x^2 + y^2)$
$g(t,x,y) = \dfrac{1}{4}y$, $h(x) = 5x$
$p(x) = e^{-2t}$, $\displaystyle\int_0^t q(s,y(s))ds = \int_0^t e^{-s}y(s)ds = x(t)$ and $\tau = 1$.




Figure 1: Stability of solutions for the nonlinear delay integro-differential equation (12)

Thus, by using Maple 2015 we have been able to establish the asymptotic stability of solutions for the second order nonlinear delay integro-differential equation (12) which is a special case of (3). This improve on the results achieve by Zhao and Meng [28].

Acknowledgement. The authors will like to appreciate the reviewers for their helpful suggestions and comments which have improved the quality of this paper.

Competing interests: The manuscript was read and approved by all the authors. They therefore declare that there is no conflicts of interest.

Funding: The authors received no financial support for the research, authorship, and/or publication of this article.

References

[1] Adams D. O. & Olutimo A. L. (2019). Some results on the boundedness of solutions of a certain third order non-autonomous differential equations with delay. *Advanced Studies in Contemporary Mathematics*. **29** (2), 237-249.

[2] Adams D. O. & Omeike M. O. (2021). Stability of solutions of certain third order non-autonomous ordinary differential equations. *Afrika Matematika*. **32** (1), 131-138. ISSN 1012-9402.

[3] Adeyanju A. A., Ademola A. T. & Ogundare B. S. (2021). On stability, boundedness and integrability of solutions of certain second order integro-differential equations with delay. *Sarajevo Journal of Mathematics*. **17** (30)1, 61-77.

[4] Ahmad S. & Stamova I. M. (Editors) (2013). *Lotka-Volterra and related systems. Recent developments in population dynamics. De Gruyter Series in mathematics and life sciences.* **2**. Berlin: De Gruyter.

[5] Berezansky L. & Domoshnitsky A. (2019). On stability of a second order integro-differential equation. *Nonlinear Dynamics and Systems Theory.* **19** (1-SI), 117-123.

[6] Burton T. A. (2005). *Volterra integral and differential equations.* Second edition Mathematics in Science and Engineering. **202**. Amsterdam: Elsevier B. V.

[7] Burton T. A. & Mahfoud W. E. (1984). *Instability and stability in Volterra equations. Trends in theory and practice of nonlinear differential equations.* (**Arlington, TX, 1982**), 99–104. Lecture Notes in Pure and Appl. Math. **90**, Dekker, New York.

[8] Corduneanu C. (1977). *Principles of differential and integral equations.* Second edition. Bronx (NY): Chelsea Publishing.

[9] Corduneanu C. & Sandberg I. W. (Editors) (2000). *Volterra equations and applications.* Papers from the Volterra centennial symposium held at the University of Texas, Arlington, TX, May 23–25, 1996. Stability and control: theory, methods and applications. **10**. Amsterdam: Gordon and Breach Science.

[10] Crisci M. R., Kolmanovskii V., Russo E. & Vecchio A. (1995). Stability of continuous and discrete Volterra integro-differential equations by Liapunov approach. *J. Integral Equations Appl.* **7** (4), 393-411.

[11] Da Prato G. & Iannelli M. (Editors) (1989). *Volterra integrodifferential equations in Banach spaces and applications.* Papers from the conference held in Trento, February 2–7, 1987. Pitman research notes in mathematics series, 190. Harlow: Longman Scientific and Technical; copublished in the United States with John Wiley and Sons, New York.

[12] Drozdov A. (1993). Stability of viscoelastic shells under periodic and stochastic loading. *Mech. Res. Comm.* **20** (6), 481-486.

[13] Goltser Y. & Litsyn E. (2008). Non-linear Volterra IDE, infinite systems and normal forms of ODE. *Nonlinear Anal.* **68** (6), 1553-1569.

[14] Gripenberg G., Londen S. Q. & Staffans O. (1990). *Volterra integral and functional equations.* encyclopedia of mathematics and its applications. **34**. Cambridge: Cambridge University Press.

[15] Lakshmikantham V. & Mohana Rao M. R. (1995). *Theory of integrodifferential equations.* Stability and control: theory, methods and applications. **1**. Lausanne: Gordon and Breach Science.

[16] Mahdavi M. (2004). Asymptotic behavior in some classes of functional differential equations. . *Non-linear Dyn. Syst. Theory.* **4** (1), 51-57.

[17] Miller R. K. (1971). Asymptotic stability properties of linear Volterra integrodifferential equations. *J. Differ. Equ.* **10**, 485-506.

[18] Morchalo J. (1991). Stability theory for Volterra equations.. *Zb. Rad. Prirod. Mat. Fak. Ser. Mat.* **21** (2), 175-183.

[19] Peschel M. & Mende W. (1986). *The predator-prey model: do we live in a Volterra world Vienna.* Springer-Verlag.

[20] Omeike O. O., Olutimo A. L. & Oyetunde O. O. (2012). The boundedness of solutions of certain nonlinear third order ordinary differential equations. *Journal of the Nigerian Mathematical Society.*. **13**, 49-54.

[21] Pi D. (2014). Stability conditions of second order integrodifferential equations with variable delay. *Abstract and Applied Analysis.* **2014**, Article ID 371639, 11 pages. Hindawi Publishing Corporation.

[22] Raffoul Y. (2007). Construction of Lyapunov functionals in functional differential equations with applications to exponential stability in Volterra integrodifferential equations. *Aust. J. Math. Anal. Appl.*. **4** (2), Art. 9, 13 pp.
[23] Staffans O. J. (1988). A direct Lyapunov approach to Volterra integro-differential equations. *SIAM J. Math. Anal.*. **19** (4), 879–901.
[24] Tunç C. (2016). New stability and boundedness results to Volterra integrodifferential equations with delay. *J. Egypt Math. Soc.* **24** (2), 210-213.
[25] Tunç C. & Tunç O. (2018). On behaviours of functional Volterra integro-differential equations with multiple tim lags. *Journal of Taibah University for Science*. **12** (2), 173-179.
[26] Wazwaz A. M. (2011). *Linear and nonlinear integral equations. Methods and applications*. Beijing/Heidelberg: Higher Education Press/Springer.
[27] Yeniçerioğlu A. F. (2008). Stability properties of second order delay integro-differential equations. *Computers and Mathematics with Applications*. **56**, 3109-3117.
[28] Zhao J. & Meng F. 2018. Stability analysis of solutions for a kind of integro-differential equations with delay. *Mathematical Problems in Engineering*. **2018**, Article ID 9519020, 6 pages. Hindawi.